

AU-AIU4 829 CARNEGIE-MELLON UNIV PITTSBURGH PA MANAGEMENT SCIENC--ETC F/6 12/1  
OPTIMAL INTEGER AND FRACTIONAL COVERS: A SHARP BOUND ON THEIR R--ETC(U)  
MAY 81 E BALAS N00014-75-C-0621  
UNCLASSIFIED MSRR-475 NL

1 of 1  
ADA  
000000




END  
DATE  
FILMED  
10-81  
DTIC

LEVEL



AD A104829




# Carnegie-Mellon University

PITTSBURGH, PENNSYLVANIA 15213

## GRADUATE SCHOOL OF INDUSTRIAL ADMINISTRATION

WILLIAM LARIMER MELLON, FOUNDER



DTIC FILE COPY

SEP 30 1981

A

This document is approved  
for public release and its  
distribution is unlimited

W.P.# 53-80-81

Management Sciences Research Report No. 475

OPTIMAL INTEGER AND FRACTIONAL

COVERS: A SHARP BOUND ON THEIR RATIO

by

Egon Balas

May 1981

The research underlying this report was supported by Grant ECS-7902506 of the National Science Foundation and Contract N00014-75-C-0621 NR 047-048 with the U.S. Office of Naval Research. It was also supported by the Alexander von Humboldt Foundation of the Federal Republic of Germany through a Senior U.S. Scientist Award. Reproduction in whole or in part is permitted for any purpose of the U.S. Government.

Management Science Research Group  
Graduate School of Industrial Administration  
Carnegie-Mellon University  
Pittsburgh, Pennsylvania 15213

\*Issued simultaneously as Report 81-13 of the Mathematisches Institut, University of Cologne.

This document has been approved  
for publication and its  
distribution is unlimited.

Abstract

The ratio of the values of optimal integer and fractional solutions to a set covering problem was shown by Johnson [5] and Lovász [6] to be bounded by  $B(d) = 1 + \frac{1}{2}d$ , where  $d$  is the largest column sum. We show that if  $n$  is the number of variables,  $B(n) = \frac{1}{n} \left\lfloor \frac{n+1}{2} \right\rfloor \left\lceil \frac{n+1}{2} \right\rceil$  is a best possible bound on this ratio. Furthermore, for every  $n \geq 20$  there are problems for which  $B(n) \leq \frac{1}{2.5} B(d)$ .

A

OPTIMAL INTEGER AND FRACTIONAL  
COVERS: A SHARP BOUND ON THEIR RATIO

by

Egon Balas

The simple (unweighted) set covering problem is

$$(C) \quad z_C = \min\{e_n x \mid Ax \geq e_m, x \text{ binary}\},$$

where  $A$  is an  $m \times n$  0-1 matrix and for  $k = m, n$ ,  $e_k$  is the  $k$ -vector whose components are all equal to 1, while  $x$  is an  $n$ -vector of variables.

If the 0-1 condition on the variables is relaxed to nonnegativity, we obtain the continuous or fractional set covering problem

$$(F) \quad z_F = \min\{e_n x \mid Ax \geq e_m, x \geq 0\}.$$

A vector  $x$  that satisfies the constraints of (C) (of (F)) will be called a cover (fractional cover).

The set covering problem is known to be NP-complete. One of the best known procedures for finding a cover that approximates the optimum is the greedy heuristic, which consists of a sequence of steps, each of which assigns the value 1 to a variable whose column covers a maximal number of additional rows. The worst case behavior of the greedy heuristic for the (unweighted) set covering problem was shown by Johnson [5] and Lovász [6] to be given by the relation

$$(1) \quad \frac{z_C}{z_F} \leq H(d) \quad (< 1 + \ln d),$$

where  $z_G$  is the value of a cover obtained by the greedy heuristic,

$$d = \max_{j \in \{1, \dots, n\}} \sum_{i=1}^m a_{ij},$$

and

$$H(d) = \sum_{j=1}^d \frac{1}{j}.$$

Thus the ratio between the value of a "greedy" cover and that of an optimal fractional cover increases at most with the logarithm of the largest column sum.

Chvátal [2] has shown that the worst case bound given by (1) is also valid for the greedy heuristic when applied to the weighted set covering problem with arbitrary but positive cost coefficients  $c_j$ ,  $j = 1, \dots, n$ . If  $k_{jt}$  represents the number of new rows covered by column  $j$  at step  $t$ , the greedy heuristic for the weighted set covering problem assigns the value 1 at step  $t$  to a variable  $x_j$  whose choice maximizes  $k_{jt}/c_j$ . Furthermore, Ho [3] has shown that the bound given by (1) is best possible for any (weighted) set covering heuristic that assigns the value 1 at step  $t$  to a variable  $x_j$  whose choice maximizes some arbitrary function  $f(c_j, k_{jt})$ .

Another class of heuristics, which uses information (reduced costs) obtained from a (not necessarily optimal) solution to the dual linear program, has consistently outperformed in empirical tests the greedy heuristic and its above mentioned generalizations (see Balas and Ho [1]), but no worst case bound better than (or comparable to) (1) is known for it (see Hochbaum [4] for a discussion of bounds for this heuristic).

Since  $z_G \geq z_C \geq z_F$ , the relation (1) implies of course both

$$(2) \quad \frac{z_G}{z_C} \leq H(d)$$

and

$$(3) \quad \frac{z_C}{z_F} \leq H(d) .$$

However, while  $H(d)$  is a best possible bound for both  $z_G/z_F$  and  $z_G/z_C$ , it was until recently an open question whether it is also a best possible bound for  $z_C/z_F$ , since no better bound than  $H(d)$  was known for this latter ratio.

In this paper we give a best possible bound on the value of  $z_C/z_F$  for unweighted set covering problems, as a function of the number  $n$  of columns, for an arbitrary number of rows. For every value of  $n$ , there are problems for which this bound has a value of approximately  $\frac{1}{2.5} H(d)$ .

For an arbitrary 0-1 matrix  $A$ , we will denote by  $z_{C(A)}$  and  $z_{F(A)}$  the value of an optimal solution to the (unweighted) set covering problem defined by  $A$ , and to the fractional set covering problem defined by  $A$ , respectively.

Let  $\mathcal{A}^n$  denote the class of 0-1 matrices with at most  $n$  columns, and let

$$\mathcal{A}^n(p) = \{A \in \mathcal{A}^n \mid z_{C(A)} = p\} .$$

**Theorem 1.** For any positive integer  $n$  and any  $p \in \{1, \dots, n\}$ ,

$$(4) \quad \min_{A \in \mathcal{A}^n(p)} z_{F(A)} = \frac{n}{n-p+1} ,$$

and the minimum in (4) is attained for the  $\binom{n}{k} \times n$  matrix  $A^*$  whose rows are all the distinct 0-1  $n$ -vectors with exactly  $n-p+1$  components equal to 1.

**Proof.** We first show that  $A^* \in \mathcal{A}^n(p)$ .  $A^*$  has  $n$  columns by assumption.

Any binary  $n$ -vector  $x$  having at least  $p$  components equal to 1 satisfies  $A^*x \geq e_q$ , where  $q = \binom{n}{k}$ , since no row of  $A^*$  has more than  $p-1$  entries equal to 0. Further, every binary  $n$ -vector  $\bar{x}$  with at most  $p-1$  components equal to 1 violates the

inequality corresponding to that particular row of  $A^*$ , whose  $p-1$  entries equal to 0 include those positions where  $\bar{x}_j = 0$ . Thus  $z_{C(A^*)} = p$ , i.e.,  $A^* \in \mathcal{A}^n(p)$ .

Next we show that  $z_{F(A^*)} = n/(n-p+1)$ . Let  $k = n-p+1$ , and let  $\tilde{x}$  be defined by  $\tilde{x}_j = 1/k$ ,  $j = 1, \dots, n$ . Let  $B$  be any  $n \times n$  nonsingular submatrix of  $A^*$ , such that every column of  $B$  has exactly  $k$  entries equal to 1. The definition of  $A^*$  guarantees the existence of  $B$ . Now let  $\tilde{u}$  be the  $q$ -vector defined by  $\tilde{u}_i = 1/k$  if the  $i^{\text{th}}$  row of  $A^*$  is a row of  $B$ ,  $\tilde{u}_i = 0$  otherwise. Then  $\tilde{x}$  and  $\tilde{u}$  are feasible solutions to the linear program  $\min\{e_n x \mid A^* x \geq e_q, x \geq 0\}$  and its dual, respectively, with value  $e_n \tilde{x} = e_q \tilde{u} = n/k$ . Hence  $\tilde{x}$  is an optimal fractional cover, and  $z_{F(A^*)} = e_n \tilde{x} = n/(n-p+1)$ .

Finally, we show that  $A^*$  minimizes  $z_{F(A)}$  over  $\mathcal{A}^n(p)$ . Assume this to be false, and let  $A^0$  be a matrix that minimizes  $z_{F(A)}$  over  $\mathcal{A}^n(p)$ , with  $z_{F(A^0)} < z_{F(A^*)}$ . Also, let  $A^* = (a_{ij}^*)$ ,  $A^0 = (a_{ij}^0)$ . W.l.o.g., we may assume that  $A^0$  has  $n$  columns, since adding columns whose entries are all equal to 0 does not change either the integer or the fractional optimum. For every  $S \subset \{1, \dots, n\}$  such that  $|S| = p-1$ ,  $A^0$  has a row  $i$  such that  $a_{ij}^0 = 0$ ,  $\forall j \in S$ ; or else  $\hat{x}$  defined by  $\hat{x}_j = 1$ ,  $j \in S$ ,  $\hat{x}_j = 0$ ,  $j \notin S$ , would be a cover with value  $p-1$ , contrary to the assumption that  $A^0 \in \mathcal{A}^n(p)$ . Hence for every row  $i$  of  $A^*$ ,  $A^0$  has a row  $k$  such that  $a_{kj}^0 \leq a_{ij}^*$ ,  $j = 1, \dots, n$ . But then  $x \geq 0$ ,  $A^0 x \geq e_r$  implies  $A^* x \geq e_q$  (where  $r$  is the number of rows of  $A^0$ ), hence  $z_{F(A^*)} \leq z_{F(A^0)}$ , a contradiction. ||

**Theorem 2.** For any  $A \in \mathcal{A}^n$ ,

$$(5) \quad \frac{z_C(A)}{z_F(A)} \leq \frac{1}{n} \left\lfloor \frac{n+1}{2} \right\rfloor \left\lceil \frac{n+1}{2} \right\rceil,$$

and this is a best possible bound.



Proof. For fixed  $p \in \{1, \dots, n\}$ , from Theorem 1

$$(6) \quad \max_{A \in \mathcal{A}^n(p)} \frac{z_{C(A)}}{z_{F(A)}} = \frac{p}{n} (n-p+1).$$

If  $p$  is allowed to vary continuously in the interval  $[1, n]$ , the right hand side of (6) is concave and attains its maximum for  $p = (n+1)/2$ . Since  $p$  has to be integer, the maximum is attained either for  $p = \lfloor \frac{n+1}{2} \rfloor$ , or for  $p = \lceil \frac{n+1}{2} \rceil$ ; namely,

$$\begin{aligned} \max_{A \in \mathcal{A}^n} \frac{z_{C(A)}}{z_{F(A)}} &= \max \left\{ \frac{1}{n} \lfloor \frac{n+1}{2} \rfloor \left( n - \lfloor \frac{n+1}{2} \rfloor + 1 \right), \frac{1}{n} \lceil \frac{n+1}{2} \rceil \left( n - \lceil \frac{n+1}{2} \rceil + 1 \right) \right\} \\ &= \frac{1}{n} \lfloor \frac{n+1}{2} \rfloor \lceil \frac{n+1}{2} \rceil. \end{aligned}$$

Another expression for the above bound is given by

$$(7) \quad \frac{1}{n} \lfloor \frac{n+1}{2} \rfloor \lceil \frac{n+1}{2} \rceil = \begin{cases} \frac{n}{4} + \frac{1}{2} & \text{if } n \text{ is even} \\ \frac{n}{4} + \frac{1}{2} + \frac{1}{4n} & \text{if } n \text{ is odd.} \end{cases}$$

Thus, the  $n$  variables set covering problem for which the ratio  $z_{C(A)}/z_{F(A)}$  attains its maximum, is the one whose coefficient matrix has exactly  $\lfloor \frac{n+1}{2} \rfloor$  1's in every row, and contains as a row every binary  $n$ -vector with  $\lfloor \frac{n+1}{2} \rfloor$  components equal to 1. For this problem,  $z_{C(A)} = \lceil \frac{n+1}{2} \rceil$  and  $z_{F(A)} = \frac{2n}{n+2-\delta}$ , where  $\delta = 0$  if  $n$  is even and  $\delta = 1$  if  $n$  is odd.

Before concluding our paper, we compare the bound on  $z_{C(A)}/z_{F(A)}$  given in Theorem 2, with the bound on  $z_{G(A)}/z_{F(A)}$  given by (1). To do this, we note that when we consider the bound  $H(d)$  given by (1) for all set covering problems defined by matrices  $A \in \mathcal{A}^n$ , the largest  $d$  that can occur (provided  $A$  has no

componentwise equal rows), happens to occur for the matrix  $A^*$  having as rows all possible 0-1  $n$ -vectors with exactly  $\lfloor \frac{n+1}{2} \rfloor$  components equal to 1. For this matrix, we denote  $d(A^*) = d^*$ , and we have

$$d^* = \binom{n-1}{\lfloor \frac{n+1}{2} \rfloor - 1} = \binom{n-1}{\lfloor \frac{n-1}{2} \rfloor}.$$

We want to assess the value of the ratio

$$(8) \quad R = \frac{1 + 2n d^*}{\lfloor \frac{n+1}{2} \rfloor \lfloor \frac{n+1}{2} \rfloor}$$

Theorem 3. For  $n \geq 2$ ,

$$(9) \quad R > 4 \frac{n-1}{n+1} 2n \left( 2 \frac{n-1}{n} \right).$$

Proof. From (8), we have

$$(10) \quad R = \frac{n}{\lfloor \frac{n+1}{2} \rfloor \lfloor \frac{n+1}{2} \rfloor} \left[ 1 + 2n \binom{n-1}{\lfloor \frac{n-1}{2} \rfloor} \right].$$

Using Stirling's formula as refined by Robbins,

$$q^q e^{-q} (2\pi q)^{1/2} e^{1/(12q+1)} < q! < q^q e^{-q} (2\pi q)^{1/2} e^{1/12q},$$

we have

$$\begin{aligned}
\binom{n-1}{\lfloor \frac{n-1}{2} \rfloor} &= \frac{(n-1)!}{\lfloor \frac{n-1}{2} \rfloor! \lceil \frac{n-1}{2} \rceil!} \\
&> \frac{(n-1)^{n-1} \cdot e^{1-n} \cdot [2\pi(n-1)]^{1/2} \cdot e^{\alpha}}{\lfloor \frac{n-1}{2} \rfloor! \lceil \frac{n-1}{2} \rceil! \cdot e^{-\lfloor \frac{n-1}{2} \rfloor - \lceil \frac{n-1}{2} \rceil} \cdot (4\pi^2 \lfloor \frac{n-1}{2} \rfloor \lceil \frac{n-1}{2} \rceil)^{1/2} \cdot e^{\beta+\gamma}} \\
&= \left( \frac{n-1}{\lfloor \frac{n-1}{2} \rfloor} \right)^{\lfloor \frac{n-1}{2} \rfloor} \cdot \left( \frac{n-1}{\lceil \frac{n-1}{2} \rceil} \right)^{\lceil \frac{n-1}{2} \rceil} \cdot \left( \frac{n-1}{2\pi \lfloor \frac{n-1}{2} \rfloor \lceil \frac{n-1}{2} \rceil} \right)^{1/2} \cdot e^{\alpha-\beta-\gamma}
\end{aligned}$$

where

$$\alpha = \frac{1}{12(n-1)+1}, \quad \beta = \frac{1}{12 \lfloor \frac{n-1}{2} \rfloor}, \quad \gamma = \frac{1}{12 \lceil \frac{n-1}{2} \rceil}.$$

Thus

$$\ell n \binom{n-1}{\lfloor \frac{n-1}{2} \rfloor} > \lfloor \frac{n-1}{2} \rfloor \ell n \frac{n-1}{\lfloor \frac{n-1}{2} \rfloor} + \lceil \frac{n-1}{2} \rceil \ell n \frac{n-1}{\lceil \frac{n-1}{2} \rceil} + \frac{1}{2} \ell n \frac{n-1}{2\pi \lfloor \frac{n-1}{2} \rfloor \lceil \frac{n-1}{2} \rceil} + \alpha - \beta - \gamma,$$

and therefore, using (10),

$$R > \frac{n \lfloor \frac{n-1}{2} \rfloor}{\lfloor \frac{n+1}{2} \rfloor \lceil \frac{n+1}{2} \rceil} \ell n \frac{n-1}{\lfloor \frac{n-1}{2} \rfloor} + \frac{n \lceil \frac{n-1}{2} \rceil}{\lfloor \frac{n+1}{2} \rfloor \lceil \frac{n+1}{2} \rceil} \ell n \frac{n-1}{\lceil \frac{n-1}{2} \rceil} + \frac{n}{\lfloor \frac{n+1}{2} \rfloor \lceil \frac{n+1}{2} \rceil} \delta,$$

where

$$\delta = 1 + \frac{1}{2} \ell n \frac{2}{\pi(n-1)} + \alpha - \beta - \gamma,$$

and we have used the fact that  $n(n-2) < (n-1)^2$  for  $n \geq 2$ .

Using  $\left\lfloor \frac{n-1}{2} \right\rfloor + \left\lceil \frac{n-1}{2} \right\rceil = n-1$  and  $2n \left\lfloor \frac{n-1}{2} \right\rfloor \geq 2n \left\lceil \frac{n-1}{2} \right\rceil$ ,

we obtain

$$(11) \quad R > \frac{(n+1)(n-1)}{\left\lfloor \frac{n+1}{2} \right\rfloor \left\lceil \frac{n+1}{2} \right\rceil} 2n \left\lfloor \frac{n-1}{2} \right\rfloor + \frac{n}{\left\lfloor \frac{n+1}{2} \right\rfloor \left\lceil \frac{n+1}{2} \right\rceil} \left( \delta - \frac{n-1}{n} 2n \left\lceil \frac{n-1}{2} \right\rceil \right).$$

As the last term is nonnegative for  $n \geq 2$ , and

$$\left\lfloor \frac{n+1}{2} \right\rfloor \left\lceil \frac{n+1}{2} \right\rceil \leq \frac{(n+1)^2}{4}, \quad \left\lceil \frac{n-1}{2} \right\rceil \leq \frac{n}{2},$$

inequality (11) implies (9). ||

The value of the righthand side in (9) is 2.5 for  $n = 20$ , and it approaches the constant  $4/2 \sim 2.769$  as  $n$  goes to infinity. Thus for the problems for which  $d = d^*$ , the bound on  $z_G/z_F$  is about  $1/2.7$  of the bound on  $z_G/z_F$ .

#### References

- [1] E. Balas and A. Ho, "Set Covering Algorithms Using Cutting Planes, Heuristics and Subgradient Optimization." Mathematical Programming Study 12, 1980, p. 37-60.
- [2] V. Chvátal, "A Greedy Heuristic for the Set Covering Problem." Mathematics of Operations Research, 4, 1979, p. 233-235.
- [3] A. Ho, "Worst Case Analysis of a Class of Set Covering Heuristics." GSIA, Carnegie-Mellon University, June 1979.
- [4] D. Hochbaum, "Approximation Algorithms for the Weighted Set Covering and Node Cover Problems." GSIA, Carnegie-Mellon University, April 1980.
- [5] D. S. Johnson, "Approximation Algorithms for Combinatorial Problems." J. Comput. System Sci., 9, 1974, p. 256-278.
- [6] L. Lovász, "On the Ratio of Optimal Integral and Fractional Covers." Discrete Mathematics, 13, 1975, p. 383-390.

REPORT DOCUMENTATION PAGE		READ INSTRUCTIONS BEFORE COMPLETING FORM	
1. REPORT NUMBER MSRR #475		2. GOVT ACCESSION NO. 3. RECIPIENT'S CATALOG NUMBER AD-11104839	
4. TITLE (and Subtitle) Optimal Integer and Fractional Covers: A Sharp Bound on Their Ratio		5. TYPE OF REPORT & PERIOD COVERED Technical Report May 1981	
7. AUTHOR(s) Egon Balas		6. PERFORMING ORG. REPORT NUMBER MSRR 475	
8. PERFORMING ORGANIZATION NAME AND ADDRESS Graduate School of Industrial Administration Carnegie-Mellon University Pittsburgh, Pa. 15213		9. CONTRACT OR GRANT NUMBER(s) N00014-75-C-0621 NR 047-048	
11. CONTROLLING OFFICE NAME AND ADDRESS Personnel and Training Research Programs Office of Naval Research (Code 434) Arlington, Va. 22217		10. PROGRAM ELEMENT, PROJECT, TASK AREA & WORK UNIT NUMBERS	
14. MONITORING AGENCY NAME & ADDRESS (if different from Controlling Office)		12. REPORT DATE May 1981	
		13. NUMBER OF PAGES 8	
		16. SECURITY CLASS. (of this report) Unclassified	
		17. DECLASSIFICATION/DOWNGRADING SCHEDULE	
15. DISTRIBUTION STATEMENT (of this Report)			
17. DISTRIBUTION STATEMENT (of the abstract entered in Block 20, if different from Report)			
18. SUPPLEMENTARY NOTES			
19. KEY WORDS (Continue on reverse side if necessary and identify by block number) Set covering, integer programming, worst case analysis			
20. ABSTRACT (Continue on reverse side if necessary and identify by block number) The ratio of the values of optimal integer and fractional solutions to a set covering problem was shown by Johnson [5] and Lovász [6] to be bounded by $B(d) = 1 + \frac{1}{2}d$ , where $d$ is the largest column sum. We show that if $n$ is the number of variables, $B(n) = \frac{1}{n} \left\lfloor \frac{n+1}{2} \right\rfloor \left\lceil \frac{n+1}{2} \right\rceil$ is a best possible bound on this ratio. Furthermore, for every $n \geq 20$ there are problems for which $B(n) \leq \frac{1}{2.5} B(d)$ .			